

Gröbner bases of $*$ -algebras and faithful operator representations.

S. Popovych, K. Vynogradov

February 1, 2008

Abstract

We give sufficient conditions for existence of a faithful representation of a $*$ -algebra in terms of its Gröbner basis. In order to do this we propose a construction of a faithful representation. This construction applies to concrete examples: $*$ -doubles, monomial $*$ -algebras, extension of a $*$ -algebras allowing Wick ordering and others. Several examples and counterexamples are also presented.

1 Introduction

We address a fundamental question in representation theory: When can an algebra (with involution) built on generators and relations be represented by unbounded operators on Hilbert space? And when is there a faithful representation? We make a case for the usefulness of Gröbner basis techniques. Now, it has been known since the early days of the algebraic approach to quantum problems (e.g., Heisenberg's commutation relations) that there are representation theoretic dichotomies: For example, the relations of the Bosons and the Fermions display different features when represented in Hilbert space. This is the issue of unbounded operators vs bounded operators. Both examples fit the theme of the paper, viz., algebras A with involution over the complex field. In the special cases when A is assumed abelian, or if A is a C^* -algebra, we have the familiar theorems of Gelfand and Naimark, but for algebras on generators and relations, the literature so far only consists of isolated classes of examples. In particular, the question of representability by Hilbert space operators has been studied for $*$ -algebras allowing Wick ordering, monomial $*$ -algebras and other (see [10, 5, 4, 3, 6]). Lance and Tapper [10, 5] undertook an attempt to treat C^* -representability of monomial $*$ -algebras with one defining relation. However there isn't a general theory yet.

Let us fix some notations. We will denote by E a pre-Hilbert space and by H a Hilbert space. Let $L(E)$ and $L(H)$ denote the $*$ -algebras of linear operators acting on these spaces. We will study the question whether a $*$ -algebra A can be embedded as a $*$ -subalgebra in $L(E)$ or $L(H)$. Note that in the latter case elements of A are represented by bounded operators. If a $*$ -algebra A is embedded in $L(E)$ and every operator $a \in A$ is bounded

then one can extend each $a \in A$ to an operator acting on the completion H of E and thus obtain an inclusion $A \hookrightarrow L(H)$. In the general case A will be represented by unbounded operators on H such that the intersection of their domains is dense. Henceforth when unbounded operators are involved the term *representation* of an algebra means that the relations of the algebra are satisfied on the common dense invariant domain. Remind that $*$ -algebra A is called C^* -representable if there is its faithful $*$ -representation by bounded operators acting on a Hilbert space or, equivalently, A can be embedded into C^* -algebra. Thus we will decompose the question of C^* -representability into two parts: the first, of algebraic nature, is to find a faithful representation of A in a pre-Hilbert space and the second one, of analytic nature, is to find out whether this representation is in bounded operators.

The question of C^* -representability of finitely-presented $*$ -algebras has been investigated in [6]. The main novelty of the approach used in [6] was the employment of Gröbner basis technique. Here we will elaborate this approach.

In present work we find sufficient conditions for a $*$ -algebra to be faithful represented in pre-Hilbert spaces. These conditions expressed in terms of its defining relations and can be effectively verified.

For $*$ -algebra given by generators and relations if not the only then at least very natural way to prove that a homomorphism is injective is to show that some linear basis is mapped into a linear basis of the image. To construct a linear basis we use the machinery of Gröbner bases developed in [2]. A brief account of it is given in the appendix.

The main property of a $*$ -algebra which enable us to construct a faithful representation is the strongly non-expanding condition (see definition 2). But this property is hard to verify in examples. So we gave several sufficient conditions (see definition 4 and theorem 4) and apply them to many concrete examples: $*$ -doubles, monomial $*$ -algebras, extension of a $*$ -algebras with Wick ordering and others.

2 Unsinkability type restrictions on Gröbner basis of a $*$ -algebra.

We will denote by F the free associative algebra since the number of generators is not important for further considerations. In this paper we will deal exclusively with finitely-generated algebras. Let us denote by F_* the free associative algebra with generators $x_1, x_2, \dots, x_m, x_1^*, x_2^*, \dots, x_m^*$. Algebra F_* is a $*$ -algebra with involution given on generators by $(x_j)^* = x_j^*$ for all $j = 1, \dots, m$. This is just associative algebra with $2m$ generators F_{2m} with a natural involution. Note that F_* is a semigroup algebra of a semigroup W of all words in generators $x_1, x_2, \dots, x_m, x_1^*, x_2^*, \dots, x_m^*$. A set $S \subseteq F$ of defining relations of an associative algebra is called Gröbner basis if it is closed under compositions (see Appendix). A Gröbner basis of a $*$ -algebra A is a Gröbner basis of A considered as an associative algebra. We need to put some extra requirements on a Gröbner basis to make it "compatible" with involution. The main requirement we impose is a generalization of

the notion of unshrinkability of the word. A set $S \subseteq F_*$ is called *symmetric* if the ideal $\mathcal{I}$ generated by S in F_* is a $*$ -subalgebra of F_* . In particular, S is symmetric if $S^* = S$. The following definition is due to P.Tapper [10]:

Definition 1 *The word $w \in W$ is called unshrinkable if it can not be presented in the form $w = d^*du$ or $w = ud^*d$ for some nonempty word d .*

Recall that a $*$ -algebra A is called *proper* if for every element $x \in A$ condition $x^*x = 0$ implies $x = 0$. A $*$ -algebra A is called *completely proper* if $M_n(\mathbb{C}) \otimes A$ is proper for all integer n . The importance of this notion follows from the fact that any bounded unital simple $*$ -algebra is C^* -representable if and only if it is completely proper [6].

P.Tapper has conjectured that $*$ -algebra $\mathbb{C}\langle x, x^* | w = 0, w^* = 0 \rangle$ is C^* -representable if and only if word w is unshrinkable. The first author in [7] proved that monomial $*$ -algebra A is completely proper if and only if its defining relations are unshrinkable. Moreover, in this case A can be faithfully represented by operators acting on pre-Hilbert space. A much wider class of $*$ -algebras for which we will prove similar results is defined below. For the notations $u \prec w$, $R_S(w)$, BW and order on W used below we refer the reader to the appendix.

Definition 2 *A symmetric subset $S \subseteq F_*$ closed under compositions will be called non-expanding if for every $u, v, w \in BW$ such that $u \neq v$ and $ww^* \prec R_S(uv^*)$ the following inequality holds $w < \sup(u, v)$. If in addition for every word $d \in BW$ the word dd^* also belong to BW we will call S strictly non-expanding. A $*$ -algebra A is called non-expanding if it possesses a Gröbner basis GB which is non-expanding and A is strictly non-expanding if some of its Gröbner bases is strictly non-expanding.*

We will show (see theorem 2) that every strictly non-expanding algebra have a faithful representation in a pre-Hilbert space.

3 Representation construction.

In this section we will show that strictly non-expanding $*$ -algebra possesses a faithful positive functional and thus is a pre-Hilbert $*$ -algebra.

Let $G \subseteq W_n$ be a subset. An *enumeration* of G is a map $\phi : G \rightarrow \mathbb{N}$ such that $u > v$ implies $\phi(u) > \phi(v)$ which is bijection on $\mathbb{N}$ if $|G| = \infty$ and on some interval $[1, n] \subseteq \mathbb{N}$ if $|G| < \infty$.

Definition 3 *Let us define the Half-word operator $H : F_* \rightarrow F_*$ by the rule $H(uu^*) = u$ for $u \in W$ and $H(v) = 0$ if v is not positive word and extend by linearity to F_* . Let us fix a set $S \subseteq F$ closed under compositions, an enumeration $\phi : BW \rightarrow \mathbb{N}$ of the corresponding linear basis and a sequence of positive real numbers $\xi = \{a_k\}_{k \in \mathbb{N}} \subset \mathbb{R}$. Define a weight functional $T_\xi^\phi : K \rightarrow \mathbb{C}$ by putting $T_\xi^\phi(u) = a_{\phi(u)}$ for any word $u \in BW$, where K denote the linear span of BW . Let $n = |BW| \in \mathbb{N} \cup \{\infty\}$.*

Let V be a vector space over $\mathbb{C}$ with basis $\{e_k\}_{k=1}^n$. Let us define a form $\langle e_i, e_j \rangle_\xi$ on the basis elements and extend it by linearity in the first argument and conjugate-linearity in the second one by the following rules. If $i = j$ then $\langle e_i, e_i \rangle_\xi = a_i$. If $i \neq j$ then there are unique elements $u, v \in BW$ such that $i = \phi(u)$, $j = \phi(v)$. Put $\langle e_i, e_j \rangle_\xi = T_\xi^\phi \circ H \circ R_S(uv^*)$.

Theorem 1 *If S is strictly non-expanding then there exist sequence $\xi_0 = \{a_k\}_{k \in \mathbb{N}} \subset \mathbb{N}$ such that the sesquilinear form $\langle \cdot, \cdot \rangle_{\xi_0}$ is positively defined.*

Proof: That the form $\langle \cdot, \cdot \rangle_\xi$ is sesquilinear is obvious by definition. Let $g_{ij} = \langle e_i, e_j \rangle_\xi$ for $i, j \in \mathbb{N}$ and $G = (g_{ij})_{1 \leq i, j \leq \infty}$ denote the Gram matrix of the sesquilinear form. We will use Silvester's criterion and show by induction on m that a_m can be chosen such that principal minor $\Delta_m > 0$. For $m = 1$ put $a_1 = 1$ then $\Delta_1 = 1 > 0$. Let $a_1, \dots, a_{m-1}$ be already chosen such that $\Delta_1 > 0, \dots, \Delta_{m-1} > 0$.

By definition if $u \in BW$, then u^*u is also in BW . Thus by definition 3 we have $\langle e_{\phi(u)}, e_{\phi(u)} \rangle_\xi = a_{\phi(u)}$. Take some $i \leq m$ and $j \leq m$ with $i \neq j$ and find unique $u, v \in BW$ such that $i = \phi(u)$, $j = \phi(v)$. Then $uv^* = \sum_k \alpha_k w_k$ for unique $\alpha_k \in \mathbb{C}$ and $w_k \in BW$. Clearly $\langle e_{\phi(u)}, e_{\phi(u)} \rangle_\xi$ is a sum $\sum_k \alpha_k a_{\phi(h_k)}$ where the sum is taken over those k for which w_k is a positive word $w_k = h_k h_k^*$. By definition we get that $h_k < \sup(u, v)$. Hence g_{ij} is a polynomial in variables $a_1, \dots, a_{m-1}$. Now from the decomposition by the m -th row we obtain $\Delta_m = \Delta_{m-1} a_m + p_m(a_1, \dots, a_{m-1})$, where $p_m \in \mathbb{C}[a_1, \dots, a_{m-1}]$ some polynomial. Since $\Delta_{m-1} > 0$ it is clear that a_m can be chosen such that $\Delta_m > 0$. This completes inductive proof. $\triangle$

The space K is isomorphic to V via the map $u \rightarrow e_{\phi(u)}$. Thus the inner product $\langle \cdot, \cdot \rangle_\xi$ on V give rise to an inner product on K which will be denoted by the same symbol. It is routine to check that $\langle u, v \rangle_\xi = \alpha(P(u \diamond v^*))$ where $P : F \rightarrow F$ is the projection on the linear span of positive words W_+ and $\alpha : K \rightarrow \mathbb{C}$ some linear functional. Let $z \rightarrow L_z$ denote the right regular representation of $A = F/\mathcal{I}$, $L_z(f) = fz$ for any $z, f \in A$.

Theorem 2 *Let $S \subseteq F$ be strictly non-expanding and $\mathcal{I}$ the ideal generated by S in F_* . Then the right regular representation L of the $*$ -algebra $A = F_*/\mathcal{I}$ on a pre-Hilbert space $(K, \langle \cdot, \cdot \rangle_\xi)$ is a faithful $*$ -representation.*

Proof: The representation stated in the theorem is associated by the GNS construction with the positive functional $\alpha(P(\cdot))$ on A . Thus it is a $*$ -representation. Indeed, as in the GNS construction $N = \{a \in A | \alpha(P(aa^*)) = 0\}$ is a right ideal in A . We can define inner product on A/N by usual rule $\langle a + N, b + N \rangle = \alpha(P(a^*b))$. It is easy to verify that right multiplication define $*$ -representation of A on pre-Hilbert A/N . The only difference with classical GNS construction is that this representation could not be, in general, extended on completion of A/N .

Let us show that $*$ -representation is a faithful $*$ -representation. Take any $f = \sum_{1 \leq i \leq n} c_i w_i \in A$, where $c_i \in \mathbb{C}, w_i \in BW$. Without loss of generality consider w_1 to be the greatest word among w_j . Then $L_f(w_1^*)$ contains element $w_1 w_1^*$ with coefficient c_1 . Hence $L_f \neq 0$. $\triangle$

Corollary 1 *Every strictly non-expanding $*$ -algebra has a faithful $*$ -representation by unbounded operators.*

4 Sufficient conditions. Examples.

We call a subset $S \subseteq F$ *reduced* if for any $s \in S$ and any word $w \prec s$ no word $\hat{s}'$ with $s' \in S$ is contained in w as a subword. If the set S is closed under compositions then one can obtain reduced set S' closed under compositions generating the same ideal by replacing each $s \in S$ with $R_S(s)$. The following, rather technical, modification of the notion of *appropriate* $*$ -algebra from [6] is the main source of examples.

Definition 4 *A symmetric reduced subset $S \subseteq F_*$ is called strictly appropriate if it is closed under compositions and for every $s \in S$ and every word $u \prec s$ such that $|u| = \deg(s)$ the following conditions hold:*

1. *word u is unshrinkable;*
2. *if $u \neq \hat{s}$, $\hat{s} = ab$, and $u = ac$ for some words a, b, c with $a \neq \emptyset$ then for any $s_1 \in S$ such that there is word $w \prec s_1$, $w \neq \hat{s}_1$, $|w| = |\hat{s}_1|$ either word $\hat{s}_1$ does not contain u as a subword or $\hat{s}_1$ and u do not form a composition in such a way that $\hat{s}_1 = d_1 a d_2$ and $u = a d_2 d_3$ with some nonempty words d_1, d_2, d_3 .*

A $*$ -algebra A is called strictly appropriate $*$ -algebra if it possesses a strictly appropriate Gröbner basis.

The following lemma shows that strictly appropriate $*$ -algebras provide examples of non-expanding $*$ -algebras. Many concrete examples of strictly appropriate $*$ -algebras will be considered in the final section. In the following lemma for word $w \in W$ of even length $w = w_1 w_2$, $|w_1| = |w_2|$ we will denote by $H_0(w)$ the first half of w , $H_0(w) = w_1$.

Theorem 3 *Every strictly appropriate set $S \subseteq F$ is non-expanding. If in addition $S = S^*$ then S is strictly non-expanding.*

Proof: Let $u, v \in BW$ and $u \neq v$.

1. If $uv^* \in BW$ then $ww^* \prec R_S(uv^*)$ implies $ww^* = uv^*$. By lemma 2 [6] two cases are possible (1) $u = vdd^*$ and $w = vd$ or (2) $v = udd^*$ and $w = vd$ where word d is nonempty. Hence $|w| = |d| + |v| = |u| - |d^*| < |u|$ in the first case and $|w| = |d| + |u| = |v| - |d^*| < |v|$ in the second one. Thus $w < u$ or $w < v$.

2. Now let $uv^* \notin BW$. There are words $p, q \in BW$ and element $s \in S$ such that $uv^* = p\hat{s}q$. Moreover, since $u, v \in BW$ none of them can contain $\hat{s}$ as a subword. Hence $\hat{s} = ab$ with nonempty words a and b such that $u = pa$ and $v^* = bq$. Write down $s = \alpha\hat{s} + \sum_{i=1}^k w_i + f$, where $\deg(f) < \deg(s)$ and $|\hat{s}| = |w_i|$ for all $i \in \{1, \dots, k\}$. Assume that for some integer i word pw_iq belongs to BW and $pw_iq = ww^*$ for some word w . If the middle of the word pw_iq comes across w_i , in other words $\max(|p|, |q|) < |w|$ then $w_i = cd$,

$w = pc$, and $w^* = dq$ with some nonempty words c, d . Hence $pc = q^*d^*$. If $|c| \leq |d|$ then $d^* = gc$ for some word g and so $w_i = cd = cc^*g^*$ which contradicts unshrinkability of w_i . If $|c| > |d|$ then $pc = q^*d^*$ implies $c = gd^*$ for some word g and we again see that $w_i = gd^*d$ is shrinkable. Thus $\max(|p|, |q|) \geq |w|$. If $|p| > |w|$ then $|u| = |p| + |a| > |w|$, otherwise $|v| = |b| + |q| > |w|$.

In the above-mentioned cases we have had $|w| < \max(|u|, |v|)$ which is a stronger statement than that of the lemma. But on the second step of the decomposition process which we now approaching this regularity breaks down.

3. Let $uv^* = p\hat{s}q$ and $s = \alpha\hat{s} + \sum_i w_i + f$ as above and $ww^* \prec R_S(pw_iq)$. It is obvious that $w \leq \sup(u, v)$. We need to prove that $w < \sup(u, v)$. Suppose the contrary: $uu^* \prec R_S(uv^*)$ and $u > v$ or $vv^* \prec R_S(uv^*)$ and $v > u$. If $|u| \neq |v|$ then $ww^* \prec R_S(pw_iq)$ implies $|w| < \max(|u|, |v|)$ because $d < c$ for all $d \prec R_S(c)$. If $|\hat{s}| > |w_i|$ then again $|w| < \max(|u|, |v|)$. Hence we can assume that $|u| = |v|$ and $|\hat{s}| = |w_i|$. Clearly $u < v$ implies $uv^* < vv^*$. Hence relation $vv^* \prec R_S(pw_iq)$ is impossible. Thus we are left with the only possibility $u > v$ and $uu^* \prec R_S(pw_iq)$. Since $uu^* < pw_iq < uv^*$ word pw_iq begins with u . If $\hat{s} = ab$ such that $pa = u, bq = v^*$ then w_i should also begin with a . Therefore $\hat{s}$ and w_i begin with the same generator. Since $pw_iq \notin BW$ there is $s_1 = \alpha\hat{s}_1 + \sum_j \beta_j u_j + g \in S$ with $\deg(g) < \deg(s_1)$ such that $pw_iq = p_1\hat{s}_1q_1$ for some words p_1, q_1 . If we assume that for some j word $uu^* \prec R_S(p_1u_jq_1)$ then $|\hat{s}_1| = |u_j|$ and $H_0(p_1u_jq_1) = u$. Word $\hat{s}_1$ could not be a subword in the first half of the word pw_iq since $H_0(p_1u_jq_1) = H_0(pw_iq) = u$ and assuming the contrary we see that $\hat{s}_1$ and u_j are both subwords of u in the same position, hence should be equal $\hat{s}_1 = u_j$. Word $\hat{s}_1$ could not contain subword w_i because of condition 1 in the definition of strictly appropriateness. Obviously, $\hat{s}_1$ could not be a subword in q because $q \in BW$. Thus either w_j and $\hat{s}_1$ intersect or $\hat{s}_1$ and w_j intersect in such a way that $\hat{s}_1 = d_1ad_2$ and $w_j = ad_2d_3$. But this contradicts the strictly appropriateness of S . So we have proved that $w < \sup(u, v)$. The fact that for any word $g \in BW$ word gg^* lies in BW follows from lemma 2. [6]. $\triangle$

The following is a convenient simplification of the preceding theorem:

Corollary 2 *If a symmetric subset $S \subseteq F$ is closed under compositions, for every $s \in S$ and every word $u \prec s$ such that $|u| = \deg(s)$ the word u is unshrinkable and words $\hat{s}$ and u begin with different generators then S is non-expanding. If in addition $S = S^*$ then S is strictly non-expanding.*

Example 1 *Let $\mathcal{L}$ be a finite dimensional real Lie algebra with linear basis $\{e_j\}_{j=1}^n$. Then its universal enveloping algebra $U(\mathcal{L})$ (over $\mathbb{C}$) is a $*$ -algebra with involution given on generators as $e_j^* = -e_j$. We claim that this $*$ -algebra is non-expanding. Indeed $M = \{e_i e_j - e_j e_i - [e_i, e_j], i < j\}$ is a set of defining relations for $U(\mathcal{L})$ it is closed under compositions (see example in [2] or use PBW theorem). Then the set $S = \{e_j^* + e_j, 1 \leq j \leq n\} \cup M$ is also closed under compositions (we consider $e_1^* > e_2^* > \dots > e_n^* > e_1 > \dots > e_n$) since e_j^* and $e_k e_l$ do not intersect for any j, k, l . It is easy to see that S is symmetric. Thus S is non-expanding by corollary 2.*

Theorem 4 *Let $S \subseteq F_*$ be a symmetric subset of a free countably generated $*$ -algebra $F_* = F(X \cup X^*)$ such that the following conditions are satisfied:*

1. *For every $s \in S$ and every word $w \prec s$ with $|w| = \deg(s)$ is unshrinkable.*
2. *For every $s_1, s_2 \in S$ and every word $u \prec s_1$ with $|u| = \deg(s_1)$ the words u and $\hat{s}_2$ do not form a composition. Then $*$ -algebra $A = \mathbb{C}\langle X \cup X^* | S \rangle$ is non-expanding. If in addition $S = S^*$ then A is strictly non-expanding.*

Proof: It suffices to show that for any two basis words $u, v \in BW$ such that $|u| = |v|$ and $u > v$ and for any word p condition $pp^* \prec R_S(uv^*)$ implies $pp^* < uu^*$.

Indeed, if $|u| \neq |v|$ then, clearly, $|pp^*| \leq |uv^*| < \max(|uu^*|, |vv^*|)$. Hence $p < \sup(u, v)$. If $|u| = |v|$ and $v > u$ then $uv^* < vv^*$. Thus every word w such that $w \prec R_S(uv^*)$ is less than vv^* .

Since $pp^* \leq uv^*$ it remains to prove that $uu^* \not\prec R_S(uv^*)$. Assume the contrary. Then there is a sequence of words $\{q_i\}_{i=1}^n$ such that $q_1 = uv^*$, $q_n = uu^*$ and for every $1 \leq i \leq n-1$ there is $s_i \in S$ and words $c_i, d_i, u_i \in W$ such that $u_i \prec s_i$, $u_i \neq \hat{s}_i$, $|u_i| = |\hat{s}_i|$ and $q_i = c_i \hat{s}_i d_i$, $q_{i+1} = c_i u_i d_i$.

Let j be the greatest with the property that $\hat{s}_j$ intersects the middle of q_j . Such an index j exists because $j = 1$ satisfies this property and we choose within a finite set. Clearly $j < n$ since otherwise u_{n-1} would be a subword in uu^* intersecting its middle and thus would be shrinkable. Thus for every $i \in \{j+1, \dots, n-1\}$ word $\hat{s}_i$ does not intersect the middle of the word $c_{i-1} u_{i-1} d_{i-1}$. But $\hat{s}_i$ could not be situated in the first half of this word because otherwise the first half of the word q_{i+1} would be strictly less than u and consequently $q_n < uu^*$ which is a contradiction. Thus $\hat{s}_i$ is a subword in the right half of the word q_{j+1} unlike that of $q_n = uu^*$ is intersected by the unshrinkable word u_j . Thus there is $k \in \{j+1, \dots, n-1\}$ such that u_j and $\hat{s}_k$ do form a composition. This contradicts to the conditions of the theorem. $\triangle$

1. Let $S = \{w_j\}_{j \in \mathbb{R}}$ be a symmetric set consisting of unshrinkable words. Since compositions of any two words are always zero this set is closed under compositions. The other conditions in the definition of strictly non-expanding set is obvious. Thus $*$ -algebra

$$\mathbb{C}\langle x_1, \dots, x_n, x_1^*, \dots, x_n^* | w_j, j \in \mathbb{R} \rangle$$

has a faithful $*$ -representation by unbounded operators.

2. Let us consider in more detail the simplest example of monomial $*$ -algebras $\mathfrak{A}_{x^2} = \mathbb{C}\langle x, x^* | x^2 = 0, x^{*2} = 0 \rangle$. It was proved in [10] that $*$ -algebra $\mathbb{C}\langle x, x^* | x^p = 0, x^{*p} = 0 \rangle$ is C^* -representable for every integer p . We will show that among the representations given by theorem 2 there is a $*$ -representation in bounded operators. We believe that this is true for any monomial $*$ -algebra with unshrinkable relations but even for $\mathbb{C}\langle x, x^* | x^3 = 0, x^{*3} = 0 \rangle$ we could not make an explicit calculations as we do for $\mathfrak{A}_{x^2}$.

It can be easily verified that BW consists of words $u_k = x(x^*x)^k$, $v_k = x^*(xx^*)^k$, $a_m = (xx^*)^m$, $b_m = (x^*x)^m$ where $k \geq 0, m \geq 1$. Obviously BW_+ consists of only words a_m and b_m ($m \geq 1$). If $z, w \in BW$ then $zw^* \in W_+$ only in the following cases

1.) $z = u_k, w = u_t$; 2.) $z = v_k, w = v_t$; 3.) $z = a_m, w = a_n$; 4.) $z = b_m, w = b_n$. Moreover,

$$u_k u_t^* = a_{k+t+1}, v_k v_t^* = b_{k+t+1}, a_m a_n^* = a_{n+m}, b_m b_n^* = b_{n+m}.$$

Consider the following ordering

$$u_0 < u_1 < \dots < a_1 < a_2 < \dots < v_0 < v_1 < \dots < b_1 < b_2 < \dots$$

Denote $\alpha(a_m) = \alpha_m$, $\alpha(b_m) = \beta_m$ then the Gram matrix of the inner product defined in theorem 1 is block diagonal

$$\begin{pmatrix} A & 0 & 0 & 0 \\ 0 & A' & 0 & 0 \\ 0 & 0 & B & 0 \\ 0 & 0 & 0 & B' \end{pmatrix}.$$

Where A, A', B, B' are Gankel matrices $A = (a_{ij})_{ij}$, $a_{ij} = \alpha_{i+j-1}$, $A' = (a'_{ij})_{ij}$, $a'_{ij} = \alpha_{i+j}$, $B = (b_{ij})_{ij}$, $b_{ij} = \beta_{i+j-1}$, $B' = (b'_{ij})_{ij}$, $b'_{ij} = \beta_{i+j}$. Note that Y' obtained from Y by cancelling out the first column (here Y stands for A or B).

Thus the question of positivity of the form $\langle \cdot, \cdot \rangle$ is reduced to the question of simultaneous positivity of two Gankel matrices C and C' where the second is obtained by cancelling out the first column. We will show that such matrices A, A', B, B' could be chosen to be positive and such that $B = A$ and that the representation in theorem 2 is in bounded operators.

Let $f : [0, 1] \rightarrow [0, 1]$ be continuous function $f(x) > 0$ for all $x \in [0, 1]$. Let

$$\alpha_m = \int_0^1 t^{m+1} f(t) dt$$

be the moments of the measure with density $f(t)$. It is well known that then moment matrix $A = (a_{ij})_{i,j=1}^n$ (where $a_{ij} = \alpha_{i+j-1}$) is positively defined. But then A' is the moment matrix of the measure with density $tf(t)$ and thus is also positively-defined. We can put $B = A$.

To prove that the representation is in bounded operators we need only to verify that the multiplication L_x by generator x is a bounded operator. Obviously, $xu_k = 0$ and $xa_m = 0$ for all $k \geq 0$ and $m \geq 1$. Compute $\|xv_k\|^2 = \langle a_{k+1}, a_{k+1} \rangle = \alpha_{2(k+1)}$, $\|v_k\|^2 = \alpha(b_{2k+1}) = \beta_{2k+1} = \alpha_{2k+1}$. Analogously, $\|xb_k\|^2 = \alpha_{2k+1}$ and $\|b_k\|^2 = \alpha_{2k}$. Thus L_x is bounded if there is a constant $c \geq 0$ such that

$$\alpha_{2(k+1)} \leq c\alpha_{2k-1}, \quad \alpha_{2k+1} \leq c\alpha_{2k}$$

for all $k \geq 1$. But

$$\alpha_{2k} = \int_0^1 t^{2k+1} f(t) dt \leq \int_0^1 t^{2k} f(t) dt = \alpha_{2k-1}$$

and

$$\alpha_{2k+1} = \int_0^1 t^{2k+2} f(t) dt \leq \int_0^1 t^{2k+1} f(t) dt = \alpha_{2k}.$$

Thus $||L_x|| \leq 1$. This proves that $\mathfrak{A}_{x,2}$ is C^* -representable.

3. The $*$ -algebra given by the generators and relations:

$$\mathbb{C}\langle a_k, k = 1, n | a_i^* a_j = \sum_{k \neq l} T_{ij}^{kl} a_l a_k^*; i \neq j \rangle,$$

with matrix of coefficient satisfying $T_{ij}^{kl} = \bar{T}_{ji}^{lk}$ is strictly appropriate by corollary 2. since no two elements from defining relations form a composition and the greatest word of any relation begins with some a_j and all other words begin with some a_k^* . Hence it has a faithful unbounded representation. Note that with additional relations $a_i^* a_i = 1 + \sum_{k,l} T_{ii}^{kl} a_l a_k^*$ we obtain so called Wick's algebras.

4. If $S \subset \mathbb{C}W(x_1, \dots, x_n)$ is closed under compositions then the $*$ -algebra

$$A = \mathbb{C}\langle x_1, \dots, x_n, x_1^*, \dots, x_n^* | S \cup S^* \rangle$$

(sometimes called $*$ -double algebra) is non-expanding (see corollary 3).

If S satisfies additionally the property that the greatest word of any relation begins with generator different from the beginnings of other longest words of this relation then A is strictly appropriate since $S \cup S^*$ is closed under compositions. In particular, if B is finite dimensional associative algebra then its table of multiplication form a set of relations S with the greatest words of length 2 and others of length 1. Thus $*$ -algebra A which $*$ -double of B . In other words, A is a free product $B_1 * B_2$, where $B_1 \simeq B_2 \simeq B$ and involution is given on the generators $b^* = \overline{\phi(b)}$ for any $b \in B_1$ and $c^* = \overline{\phi^{-1}(c)}$ for any $c \in B_2$ with $\phi : B_1 \rightarrow B_2$ being any isomorphism.

Theorem 5 *Let $S \subset F$ be a symmetric subset of a free $*$ -algebra in generators $x_1, \dots, x_n$ and $x_1^*, \dots, x_n^*$ closed under compositions such that for any $s \in S$ the following properties holds:*

1. $\hat{s} \in G = W(x_1, \dots, x_n)$ or $\hat{s} \in G^* = W(x_1^*, \dots, x_n^*)$.
2. for any $u \prec s$ such that $|u| = |\hat{s}|$ words u and $\hat{s}$ both lie in the same semigroup G or G^* .

Then S is non-expanding.

Proof: If some word $w \in W$ ($w = y_{k_1} \dots y_{k_t}$ where $y_{k_r} \in X \cup X^*$ are generators) contains subword $\hat{s}$ with $s \in S$. Then $w = p\hat{s}q$ for some words p and q in W . After substitution $w \rightarrow p\bar{s}q = \sum_i \alpha_i p w_i q$ we see that all words w_i such that $|w_i| = |\hat{s}|$ lie in the same semigroup either in G or in G^* . Since decomposition $R_S(w) = \sum_j \beta_j u_j$, $u_j \in BW$, $u_j = z_{i_1} \dots z_{i_r}$, $z_{i_r} \in X \cup X^*$ can be obtained by several steps of substitution considered above we see that for any j such that $|u_j| = |w|$ for all r both generators z_{i_r} and y_{k_r} lie in the same set X or X^* .

Let $u, v \in BW$, $u \neq v$. Let word w be such that $ww^* \prec R_S(uv^*)$. If $|u| \neq |v|$ then $|w| < \max(|u|, |v|)$ hence $w < \sup(u, v)$. So assume that $|u| = |v|$ then $ww^* \leq uv^*$

implies that $w \leq u$. Let $w = u$. Write down $u = z_1 \dots z_k, z_r \in X \cup X^*$. Without loss of generality assume $z_k \in X$. Then $uu^* = z_1 \dots z_k z_k^* \dots z_1^*$ by the first part of the proof u ends with generator from X whereas v^* begins with generator from X^* . Thus $uv^* \in BW$ and $R_S(uv^*) = uv^*$. Clearly, $uv^* = uu^*$ implies $u = v$. Obtained contradiction proves that if $ww^* \prec R_S(uv^*)$ then $w < \sup(u, v)$. Since S is appropriate for any $d \in BW$ word dd^* lies in BW by lemma 2 [6]. $\triangle$

It could be shown using Zorn's lemma that for any algebra A and any its set of generators S there is a Gröbner basis corresponding to S with any given inductive ordering of the generators. Thus we have the following corollary.

Corollary 3 *If B is a finitely generated associative algebra then its $*$ -double $A = B * B$ is strictly non-expanding $*$ -algebra. Hence A has a faithful $*$ -representation in pre-Hilbert space.*

Below we give some known examples of $*$ -doubles which have finite Gröbner bases.

5. Consider $*$ -algebra:

$$Q_{4,\alpha} = \mathbb{C}\langle q_1, \dots, q_4, q_1^*, \dots, q_4^* \mid q_j^2 = q_j, \sum_j q_j = \alpha, q_j^{*2} = q_j^*, \sum_j q_j^* = \bar{\alpha} \rangle.$$

It is a $*$ -double of the algebra

$$B_{n,\alpha} = \mathbb{C}\langle q_1, \dots, q_4 \mid q_j^2 = q_j, \sum_j q_j = \alpha \rangle$$

which has the following Gröbner basis:

$S = \{q_1q_1 - q_1, q_2q_2 - q_2, -q_3q_2 - 2q_1 - 2q_2 - 2q_3 + \alpha + 2\alpha q_1 + 2\alpha q_2 + 2\alpha q_3 - \alpha^2 - q_1q_2 - q_1q_3 - q_2q_1 - q_2q_3 - q_3q_1, q_3q_3 - q_3, -q_3q_1q_2 - 3\alpha + 5\alpha^2 - 2\alpha^3 + q_2(6 - 10\alpha + 4\alpha^2) + q_3(6 - 10\alpha + 4\alpha^2) + q_1(8 - 13\alpha + 5\alpha^2) + (3 - 2\alpha)q_1q_2 + (6 - 4\alpha)q_1q_3 + (6 - 4\alpha)q_2q_1 + (6 - 4\alpha)q_2q_3 + (3 - 2\alpha)q_3q_1 + q_1q_2q_1 + q_1q_2q_3 + q_1q_3q_1 + q_2q_1q_3 + q_2q_3q_1\}$. More about this algebra can be found in [9, 1]. Let us note that when $\alpha = 0$ $*$ -algebra $Q_{4,0} = B_{4,0} * B_{4,0}$ has only zero representation in bounded operators (see [1]). Thus for this $*$ -algebra unboundedness is essential.

6. That the generators in the previous example are idempotents is not important, we can consider other powers as well:

$$T_{3,\alpha} = \mathbb{C}\langle q_1, q_2, q_3, q_1^*, q_2^*, q_3^* \mid q_j^3 = q_j, \sum_j q_j = \alpha, q_j^{*3} = q_j^*, \sum_j q_j^* = \bar{\alpha} \rangle.$$

It is a $*$ -double of the algebra $\mathbb{C}\langle q_1, q_2, q_3 \mid q_j^3 = q_j, \sum_j q_j = \alpha \rangle$. We can find its Gröbner basis. We have the following set of relations $\{q_1^3 - q_1, q_2^3 - q_2, q_3^3 - q_3, q_1 + q_2 + q_3 - \alpha\}$. From these relations we see that this algebra is generated by q_1 and q_2 . That is why we can consider the following set of relations: $\{q_1^3 - q_1, q_2^3 - q_2, (\alpha - q_1 - q_2)^3 - (\alpha - q_1 - q_2)\}$. Consider the following order on generators $q_2 > q_1$. All relations are already normalized. The greatest words in this relations are q_1^3, q_2^3 and $q_1^2q_2$. Thus we have no reductions. The

first and third relations give two compositions. From one side they intersect by the word q_1 . And the result of this composition is $(q_1^3 - q_1)q_1q_2 - q_1^2((\alpha - q_1 - q_2)^3 - (\alpha - q_1 - q_2))$. From other side they intersect by the word q_1^2 . Result of this composition is $(q_1^3 - q_1)q_2 - q_1((\alpha - q_1 - q_2)^3 - (\alpha - q_1 - q_2))$. Another composition is formed by third and second relations. Their greatest words intersect by the word q_2 . Result of this composition is $((\alpha - q_1 - q_2)^3 - (\alpha - q_1 - q_2))q_2^2 - q_1^2(q_2^3 - q_2)$. Hence we have three new relations. After performing reductions we will have the following set of relations:

$$S = \{q_1^3 - q_1, -q_2^2q_1 + 3\alpha q_1^2 + 3\alpha q_2^2 + \alpha^3 + q_1(-1 - 3\alpha^2) + q_2(-1 - 3\alpha^2) + 3\alpha q_1q_2 - q_1q_2^2 - q_1^2q_2 + 3\alpha q_2q_1 - q_2q_1^2 - q_1q_2q_1 - q_2q_1q_2, q_2^3 - q_2, -q_2q_1q_2q_1^2 + -\alpha^3 + 9\alpha^5 - q_1^2(-3\alpha - 37\alpha^3) - q_2^2(3\alpha - 27\alpha^3) - q_2(-1 + 6\alpha^2 + 27\alpha^4) - q_1(18\alpha^2 + 30\alpha^4) - (-12\alpha - 45\alpha^3)q_1q_2 - 27\alpha^2q_1q_2^2 - (1 + 30\alpha^2)q_1^2q_2 + 9\alpha q_1^2q_2^2 - (6\alpha - 18\alpha^3)q_2q_1 - (1 + 3\alpha^2)q_2q_1^2 - (-2 + 15\alpha^2)q_1q_2q_1 + 3\alpha q_1q_2q_1^2 + 3\alpha q_1^2q_2q_1 - q_1^2q_2q_1^2 - (-1 + 9\alpha^2)q_2q_1q_2 + 6\alpha q_1q_2q_1q_2 - q_1^2q_2q_1q_2 - 3\alpha q_2q_1q_2q_1 + q_1q_2q_1q_2q_1\}$$

Some of these relations do form compositions but all of them reduce to zero. Hence it is a Gröbner basis.

5 APPENDIX: Noncommutative Gröbner bases.

For the convenience of the reader we repeat the relevant material from noncommutative Gröbner bases theory (see [11, 2]) with some easy reformulations. Let W_n denote the free $*$ -semigroup with generators $x_1, \dots, x_n$. For a word $w = x_{i_1}^{\alpha_1} \dots x_{i_k}^{\alpha_k}$ (where $i_1, i_2, \dots, i_k \in \{1, \dots, n\}$, and $\alpha_1, \dots, \alpha_k \in \mathbb{N} \cup \{0\}$) the length of w , denoted by $|w|$, is defined to be $\alpha_1 + \dots + \alpha_k$. Let $F_n = \mathbb{C}\langle x_1, \dots, x_n \rangle$ denote the free associative algebra with generators $x_1, \dots, x_n$. We will sometimes omit subscript n . Fix the linear order on W_n such that $x_1 > x_2 > \dots > x_n$, the words of the same length ordered lexicographically and the words of greater length are considered greater. Any $f \in F_n$ is a linear combination $\sum_{i=1}^k \alpha_i w_i$ (we may assume that $\alpha_i \neq 0$ for all $i \in \{1, \dots, k\}$) of words $w_1, w_2, \dots, w_k$. Let $\hat{f}$ denote the greatest of these words, say w_j . Then denote $\hat{f} - (\alpha_j)^{-1}f$ by $\bar{f}$. The degree of $f \in F_n$, denoted by $\deg(f)$, is defined to be $|\hat{f}|$. Elements of the free algebra F can be identified with functions $f : W \rightarrow \mathbb{C}$ with finite support via the map $f \rightarrow \sum_{w \in W} f(w)w$. For a word $z \in W$ and an element $f \in F$ we will write $z \prec f$ if and only if $f(z) \neq 0$.

Definition 5 We will say that two elements $f, g \in F_n$ form a composition $w \in W$ if there are words $x, z \in W$ and nonempty word $y \in W$ such that $\hat{f} = xy$, $\hat{g} = yz$ and $w = xyz$, in other words $w = \hat{f}w_1 = w_2\hat{g}$ for some nonempty words $w_1, w_2 \in W$ in which the marked words $\hat{f}$ and $\hat{g}$ "intersect" $|\hat{f}| > |w_2|$. Let us denote the result of the composition $\beta f w_1 - \alpha w_2 g$ by $(f, g)_w$, where α and β are the coefficients of the greatest words $\hat{f}$ and $\hat{g}$ in f and g respectively.

It is obvious that $(f, g)_w \prec w$. Notice that two elements f and g may form compositions in many ways and f may form composition with itself. The following definition is due to Bokut [2].

Definition 6 A subset $S \subseteq F_n$ is called closed under compositions if for any two elements $f, g \in S$ the following properties holds:

1. if $f \neq g$ then the word $\hat{f}$ is not a subword in $\hat{g}$;
2. If f and g form a composition w then there are $a_j, b_j \in W_n, f_j \in S, \alpha_j \in \mathbb{C}$ such that $(f, g)_w = \sum_{j=1}^m \alpha_j a_j f_j b_j$ and $a_j f_j b_j < w$, for $j = 1, \dots, m$.

Definition 7 A set $S \subseteq F$ is called a Gröbner basis of an ideal $\mathcal{I} \subseteq F$ if for any $f \in \mathcal{I}$ there is $s \in S$ such that $\hat{s}$ is a subword in $\hat{f}$. A Gröbner basis S of $\mathcal{I}$ is called minimal if no proper subset of S is a Gröbner basis of $\mathcal{I}$.

If S is closed under compositions then S is a minimal Gröbner basis for the ideal $\mathcal{I}$ generated by S [2]. Henceforth we will consider only minimal Gröbner bases. Thus we will say that S is a Gröbner basis of an associative algebra $A = F/\mathcal{I}$ if S is closed under composition and S generate $\mathcal{I}$ as an ideal of F . Let GB be a Gröbner basis for A , $\hat{GB}$ be the set of greatest words $\hat{s}$ of elements of $s \in GB$ and $BW(GB)$ be the subset of those words in W_n that contain no word from $\hat{GB}$ as a subword. It is a well known that $BW(GB)$ is a linear basis for A . Further on we will write simply BW since we will always deal with a fixed Gröbner basis.

Let $S \subseteq F$ be closed under compositions, $\mathcal{I}$ the ideal generated by S . Let us define the operator $R_S : F \rightarrow F$ by the following rule. For $f \in F$ there are uniquely defined coefficients $\{\alpha_i\} \subset \mathbb{C}$ and words $\{w_i\} \subset BW$ such that $f + \mathcal{I} = \sum_i \alpha_i (w_i + \mathcal{I})$ (basis decomposition in the factor algebra). Put $R_S(f) = \sum_i \alpha_i w_i$. Then R_S is a retraction on a subspace K in F spanned by BW . We can consider the space K with the new operation $f \diamond g = R_S(fg)$ for $f, g \in K$. Then $(K, +, \diamond)$ becomes an algebra isomorphic to $F/\mathcal{I}$. The decomposition of element $f = \sum_j \beta_j u_j$ where $u_j \in W$ with respect to the basis BW in the factor algebra $A/\mathcal{I}$ can be obtained by repeated application of the following procedure: if word u_j contains subword $\hat{s}$ with $s \in S$, then there are words p and q such that $u_j = p\hat{s}q$. Substitute $\bar{s}$ instead of $\hat{s}$ (we will denote this substitution by $p\hat{s}q \rightarrow p\bar{s}q$). Having done this for all j we obtain an element $\sum_j \beta'_j u'_j \in F$. Repeat this procedure for all u'_j and so on. After finite steps we obtain desired decomposition $\sum_i \alpha_i w_i$ where all $w_j \in BW$. From this follows that if $w \prec R_S(u)$ for some words w and u then $w < u$.

Acknowledgements.

The authors wish to express their thanks to their teacher Prof. Yurii Samoilenko for his kind attention and many helpful comments.

References

- [1] H. Bart, T. Erhardt, B. Sillbermann, Zero Sums of Idempotens in Banach Algebras// *Integr. Equat. Oper. Th.*, 19 (1994) 123–134.
- [2] Bokut, ‘Embeddings of rings’, *Uspehi mat. nauk* 42 No.4 (1987) 87-111. (in Russian)
- [3] Palle E. T. Jorgensen and Daniil P. Proskurin and Yurii S. Samoilenko, ‘The kernel of Fock representations of Wick algebras with braided operator of coefficients’,

- Pacific J. Math.*, 198 (2001) 109–122; arXiv:math-ph/0001011 (2001), pp. 321–329; arXiv:math.QA/0010308
- [4] P.E.T Jorgensen, L.M. Schmitt, R.F. Werner, ‘Positive representations of general commutation relations allowing wick ordering’, *Funct. Anal.* 134, No.1 (1995), 33-99; arXiv:funct-an/9312004.
 - [5] E. C. Lance, ‘Finitely-presented C^* -algebras . 1997, Operator algebras and applications’, (ed. A. Katavolos), Nato ASI Series, Ser. C: Math and Phys. Sci., Vol.495, Kluwer Acad. Publ., 1997, pp.255-266
 - [6] S. V. Popovych , ‘Conditions for embedding a $*$ -algebra into a C^* -algebra’, *Methods of Funct. Analysis and topology.* 5, No.3 (1999) 40-48.
 - [7] S. V. Popovych, ‘Monomial $*$ -Algebras and Tapper’s Conjecture’ *Methods of Funct. Analysis and topology.*, 8 No. 1 (2002) 70-75
 - [8] S. V. Popovych, ‘Realization of finitely presented $*$ -algebras by Hilbert space operators’, *Functional analysis:Proc. of Ukr. Math. Congress - 2001*, (Kyiv:Inst. of Math. of NAS of Ukraine, 2002), pp. 245-253
 - [9] V. I. Rabanovich, Yu. S. Samoilenko, ‘When a sum of projections is a multiple of identity’, *Funct. Anal. and Appl.*, 34 No.4 (2000) 91–93. (in Russian)
 - [10] P. Tapper, ‘Embedding of $*$ -algebras into C^* -algebras and C^* -ideals generated by words’, *J. Operator Theory*, 41 (1999) 351-364.
 - [11] V. A. Ufnarovsky , ‘I. Combinatorial and asimptotic methods in algebra’, *Sovrem. Probl. Matem. Fundam. Naprav.* 57 (1990) 5-177 (in Russian)